

## Quality of Life among Rural Postmenopausal Women in Bangladesh

Julliat Binta Ali<sup>1\*</sup>, Afroza Begum<sup>2</sup>, Khaled Rahman Ayon<sup>3</sup>

<sup>1</sup>MBBS, MPH (Reproductive and Child Health), Lecturer, Faridpur Medical College, Faridpur, Bangladesh.

ORCID ID: 0000-0003-1140-4388

<sup>2</sup>MBBS, MPH, Professor and Ex- Head of the Department (Reproductive and Child Health) National Institute of Preventive and Social Medicine (NIPSOM), Mohakhali, Dhaka-1212, Bangladesh. ORCID ID: 0000-0001-9547-2569

<sup>3</sup>Software Engineer. ORCID ID: 0000-0002-8765-4968

\***Corresponding Author:** Julliat Binta Ali, MBBS, MPH (Reproductive and Child Health), Lecturer, Faridpur Medical College, Faridpur, Bangladesh. ORCID ID: 0000-0003-1140-4388.

**Received date:** 20 December 2024; **Accepted date:** 22 January 2025; **Published date:** 29 January 2025

**Citation:** Binta Ali J, Begum F, Ayon KR (2025) *Quality of Life among Rural Postmenopausal Women in Bangladesh*. J Comm Med and Pub Health Rep 6(01): <https://doi.org/10.38207/JCMPHR/2025/JAN06010406>

**Copyright:** © 2025 Julliat Binta Ali. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Introduction

Thousands of years ago, the Roman philosopher Lucius Annaeus Seneca emphasized the significance of quality of life (QOL) by stating, "It is quality rather than quantity that matters" [1, 2]. A key objective of health services for all individuals in the twenty-first century is to enhance quality of life [3]. The QOL in the postmenopausal period has taken a lot of attention particularly in recent years since nearly one-third of women are living in postmenopausal age [1].

QOL has been outlined by the World Health Organization (WHO) as the 'individual's perceptions of their position in life within the context of the cultural and value systems within which they live and in relevancy their goals, expectations, standards, and considerations' [1,4,5,6,7,8,9,10]. Quality of life (QOL) in relation to health encompasses multiple dimensions, including social, physical, and mental aspects that affect everyday life [11, 12, 13]. Every woman experiences a unique impact on her QOL after menopause, shaped by these key areas [11].

Menopause is defined as the permanent absence of menstruation for 12 consecutive months or the loss of ovarian function leading to permanent amenorrhea [14, 15, 16]. Menopause is a natural phase in every woman's life, leading to a range of vasomotor, psychological, physical, and sexual symptoms [14]. There is a drop in the levels of estrogen and progesterone, the two most important hormones in the female body [17]. As a result of a lack of estrogen, a woman may experience decrease physical and mental well-being that is why a postmenopausal woman can be considered a risk population [13,18,19].

Menopause usually occurs on average between the ages of 50 and 51 and literally refers to a woman's last menstruation period; however, age 40 is used as an arbitrary cut-off point for natural menopause. Women who have not menstruated during the past 12 months are said to be at the postmenopausal stage [17].

The duration, intensity, and effects of menopausal symptoms can differ significantly among individuals and populations. For some women, the symptoms are severe enough to deeply impact their daily lives, personal well-being, social interactions, and overall quality of life [7,20].

Over 80% of women experience physical or psychological symptoms in the year leading up to menopause, with varying degrees of discomfort and disruption in their lives, ultimately resulting in a reduced quality of life [18,21]. Despite a majority of women facing multiple symptoms, the literature still presents a gap on groups of symptoms consistently occur and what these symptoms be affected by the quality of life of postmenopausal women [7,22].

Psychological, physical, vasomotor and sexual changes occurring in the postmenopausal period may disturb a woman's quality of life. Also, the quality of life of a woman in the post-menopausal age group is influenced by socio-cultural and behavioral factors compounded by their health-related perceptions and health-seeking behavior. Postmenopausal symptoms can lead to social consequences such as disruptions in women's domestic roles and economic activities, as well as harm to their psychological and emotional well-being, ultimately impacting their quality of life [3].

The majority of studies on the quality of life of postmenopausal women have been carried out in developed countries, where different socio-cultural factors may shape both the perception of quality of life and the experience of menopausal symptoms. However, there is limited data available on the quality of life of postmenopausal women in developing countries [7,18].

This study aims to highlight the extent of health-related challenges faced by postmenopausal women and examine the relationship between these issues and various socio-demographic factors. Additionally, it seeks to raise awareness among health authorities,

encouraging them to implement appropriate measures to address the needs of menopausal women.

## Methodology

This cross-sectional study was conducted among rural postmenopausal women in Faridpur district in Bangladesh from January, 2018 to December, 2018.

The study included all postmenopausal women residing in the study area who met the established inclusion and exclusion criteria.

### Inclusion criteria:

Women eligible for the study were those aged 40 to 70 years, who had not experienced menstruation in the past 12 months, and had undergone natural menopause.

### Exclusion criteria:

The women who had attained menopause surgically, who were severely ill physically and/or mentally and who were not willing to participate in the study were excluded from the study.

In this study, a total of 384 postmenopausal women were selected using a convenient sampling method. Data collection was carried out through face-to-face interviews, using a structured questionnaire.

The questionnaire was divided into two sections:

### Part 1: Socio-Demographic Characteristics

This section gathered personal information about the participants, including age, gender, religion, educational background, family structure, monthly income, and expenses. It also covered marital status, number of children, and age at menopause. Additional variables such as the husband's educational level, occupation, type of house, total family income, and family expenditures were also included.

### Part 2: MENQOL Questionnaire for Quality of Life

To assess the participants' quality of life, the MENQOL (Menopause-Specific Quality of Life) scale was used. This self-report measure evaluates the presence and severity of menopausal symptoms and how much they impact daily life. The MENQOL consists of 29 items, each corresponding to one of four domains of menopausal symptoms experienced in the past month:

- **Vasomotor** (items 1–3)
- **Psychosocial** (items 4–10)
- **Physical** (items 11–26)
- **Sexual** (items 27–29)

For each item, participants indicate whether the symptom was present, and if so, how bothersome it was on a scale from 0 (not bothersome) to 6 (extremely bothersome). The responses are

systematically converted for scoring and data analysis. Each seven-point Likert scale response is translated into an eight-point scale ranging from 1 to 8. A score of "1" means the participant did not experience the symptom in the last month, while a "2" indicates the symptom was present but not bothersome. Scores from "3" to "8" reflect increasing levels of bother, corresponding to the original 1-6 range.

The final score for each MENQOL domain is the average of the converted item scores and ranges from 1 to 8. The severity of menopausal symptoms is classified as follows:

- **Mild:** Score range 2–4
- **Moderate:** Score of 5
- **Severe:** Score range 6–8.

Data were collected, coded, tabulated, and analyzed using SPSS software (version 20, IBM Corporation, Armonk, NY, USA). Descriptive statistics were applied to calculate percentages, frequencies, means, and standard deviations. To assess statistical significance, Chi-square tests, t-tests, and ANOVA were conducted. A p-value of less than 0.05 ( $p < 0.05$ ) was considered statistically significant.

Formal ethical approval for this study was granted by the Institutional Review Board (IRB) of the National Institute of Preventive and Social Medicine (NIPSOM). Prior to data collection, informed written consent was obtained from each participant after providing a clear explanation of the study's objectives and purpose. All participants were treated with respect and equality, and their privacy and confidentiality were strictly upheld. Participation was entirely voluntary, and individuals were enrolled in the study only after signing the consent form.

## Results

The mean age of the women in the study was  $58.02 \pm 7.574$  years, with a mean menopausal age of  $47.13 \pm 4.300$  years. The vast majority of participants (98%) were followers of Islam. Most of the respondents (97.4%) were housewives, and 92.7% were illiterate. In terms of marital status, 60.9% were married, 36.2% were widowed, and 2.9% were divorced. Among the 384 women surveyed, 51% had four or more children, while 49% had between zero and three children. A significant portion (63%) lived in joint families, and 60.9% resided in Klay houses. Additionally, 86.4% of the participants' husbands were illiterate, with only basic literacy skills. In terms of occupation, the majority (60%) of their husbands were engaged in agricultural work. [Table 1]

**Table 1:** Socio-demographic characteristics of the studied participants

Characteristics	Number	Percentage (%)
<b>Age in years</b>		
<50	69	18
>=50	315	82
<b>Religion</b>		
Muslim	378	98
Hindu	6	2
<b>Level of education of the respondents</b>		
Illiterate	356	92.7
Primary	24	6.3
Secondary	4	1
<b>Occupation of the respondents</b>		
Housewife	374	97.4
Day labor	10	2.6
<b>Marital status</b>		
Married	234	60.9
Widow	139	36.2
Divorced	11	2.9
<b>Number of children of the respondents</b>		
0-3	188	49.0
4 and more	196	51.0
<b>Type of family of the respondents</b>		
Joint family	240	63
Nuclear family	144	37
<b>Type of housing of the respondents</b>		
Klay house	234	60.9
Semi Concrete house	100	26
Concrete house	50	13
<b>Education level of the respondent's husband</b>		
Illiterate	332	86.4
Primary	36	9.4
Secondary	9	2.3
Higher secondary	4	1
Honors	3	.8
<b>Occupation of the respondent's husband</b>		
Farmer	232	60
Day labor	84	22
Businessman	39	10
Retired	29	8
<b>Age of menopause of the respondents</b>		
40-45	152	39.6
46-50	144	37.5
51-55	88	22.9

**Table 2** indicated that the highest mean scores of symptoms in the vasomotor, psychosocial, physical and sexual domain were hot Flushes (7.49±1.101), experiencing poor memory (7.66±.982), aching

in muscles and joints (7.57±1.148) and involuntary urination when laughing or coughing (7.57±1.056), vaginal dryness during intercourse (7.57±1.056) respectively.

**Table 2:** Mean Scores of MENQOL items (N=384)

Symptoms	N	Mean ± SD
<b>A. Vasomotor</b>		
Hot flushes	363	7.49±1.101
Night Sweats	317	7.15±1.498
Sweating	377	6.97±1.693
<b>B. Psychosocial</b>		
Dissatisfaction with personal life	310	6.93±2.011
Feeling anxious or nervous	338	7.07±1.424
Experiencing poor memory	377	7.66±.982
Accomplishing less than I used to	383	7.52±1.249
Feeling depressed, down or blue	342	6.87±1.690
Impatience with other people	197	6.94±1.402
Willing to be alone	122	6.37±2.050
<b>C. Physical</b>		
Flatulence (Wind) or gas pain	316	7.01±1.613
Aching in muscles and joints	348	7.57±1.148
Feeling tired or worn out	373	7.17±1.256
Difficulty in sleeping	296	7.26±1.272
Aches in back of neck or head	339	7.33±1.422
Decreases in physical strength	384	7.09±1.719
Decreased stamina	383	7.07±1.720
Feeling lack of energy	384	7.10±1.746
Dry skin	383	6.29±2.504
Weight gain	119	6.51±1.991
Increased facial hair	2	6.50±.707
Changes in appearance, texture or tone of skin	384	6.35±2.513
Feeling bloated	202	6.85±1.866
Low Backache	323	7.47±1.286
Frequent urination	232	7.39±1.104
Involuntary urination when laughing or coughing	199	7.57±1.056
<b>D. Sexual</b>		
Change in sexual desire	225	7.45±1.420
Vaginal dryness during intercourse	212	7.57±1.114
Avoiding intimacy	224	7.44±1.403

**Table 3** illustrated the severity of the menopausal symptoms among the respondents. It was observed that the most severe symptoms in the vasomotor domain were hot flushes (85.7%). In the psychosocial domain, the most severe symptoms were experiencing poor memory (92.4%) and accomplishing less than they used to (91.4%). In

physical domain the most severe symptoms were feeling lack of energy (86.5%) and decrease in physical strength (86.2%). Out of 60.9% married postmenopausal women, the most severe symptoms in sexual domain were changed in sexual desire (52.6%) followed by avoiding intimacy (52.1%).

**Table 3:** Distribution of the postmenopausal women according to the severity of menopausal symptoms (N=384)

Symptoms	N	Mild		Moderate		Severe	
		n	%	n	%	n	%
<b>A. Vasomotor</b>							
Hot flushes	363	10	2.6	24	6.3	329	85.7
Night sweats	317	21	5.5	31	8.1	265	69.0
Sweating	377	40	10.4	39	10.2	298	77.6
<b>B. Psychosocial</b>							
Dissatisfaction with personal life	310	45	11.7	7	1.8	258	67.2
Feeling anxious or nervous	338	26	6.8	31	8.1	281	73.2
Experiencing poor memory	377	10	2.6	12	3.1	355	92.4
Accomplishing less than I used to	383	16	4.2	16	4.2	351	91.4
Feeling depressed, down or blue	342	34	8.9	26	6.8	282	73.4
Impatience with other people	197	15	3.9	16	4.2	166	43.2
Willing to be alone	122	21	5.5	14	3.6	87	22.7
<b>C. Physical</b>							
Flatulence (wind) or gas pain	316	31	8.1	34	8.9	251	65.4
Aching in muscles and joints	348	16	4.2	10	2.6	322	83.9
Feeling tired or worn out	373	15	3.9	29	7.6	329	85.7
Difficulty in sleeping	296	10	2.6	33	8.6	253	65.9
Aches in back of neck or head	339	22	5.7	24	6.3	293	76.3
Decrease in physical strength	384	37	9.6	16	4.2	331	86.2
Decreased stamina	383	37	9.6	16	4.2	330	85.9
Feeling lack of energy	384	39	10.2	13	3.4	332	86.5
Dry skin	383	97	25.3	4	1.0	282	73.4
Increased facial hair	2					2	.5
Weight gain	119	22	5.7	3	.8	94	24.5
Changes in appearance, texture or tone of skin	384	96	25.0	5	1.3	283	73.7
Feeling bloated	202	26	6.8	8	2.1	168	43.8
Low backache	323	19	4.9	12	3.1	292	76.0
Frequent urination	232	5	1.3	19	4.9	208	54.2
<b>D. Sexual</b>							
Change in sexual desire	225	15	3.9	8	2.1	202	52.6
Vaginal dryness during intercourse	212	8	2.1	9	2.3	195	50.8
Avoiding intimacy	224	14	3.6	10	2.6	200	52.1

**Table 4** illustrated the overall scores of menopausal quality of life for each MENQOL domain. It was observed that the highest mean score in sexual domain (7.42±1.387) followed by vasomotor (7.11±1.278)

then psychosocial domain (7.09±.961) and finally physical domain (7.00±1.079).

**Table 4:** Mean Score for each MENQOL domain (N=384)

Domain	Mean ± SD	Level of severity
Vasomotor	7.11±1.278	Severe
Psychosocial	7.09±.961	Severe
Physical	7.00±1.079	Severe
Sexual	7.42±1.387	Severe

**Table 5** highlights the comparison of MENQOL questionnaire scores across the four domains for women with varying socio-demographic characteristics. The analysis revealed that in the psychosocial domain, age and educational qualification of the respondents were significant factors. In the physical domain, the respondent's educational

qualification, their husband's educational level, and the type of house were key predictors. In the sexual domain, the respondent's occupation emerged as a predictor of better quality of life among postmenopausal women.

**Table 5:** Mean scores per domain in menopausal women according to socio-demographic characteristics

Socio-demographic characteristics	Vasomotor	Psychosocial	Physical	Sexual
Age				
<50	7.08±1.312	6.85±1.047	7.05±.901	7.28±1.441
≥50	7.12±1.273	7.15±.935	6.99±1.115	7.47±1.367
	p=.817	<b>p=.022*</b>	p=.646	p=.352
<b>Religion</b>				
Islam	7.12±1.275	7.10±.961	7.02±1.063	7.42±1.394
Hinduism	6.55±1.430	6.63±.936	6.22±1.658	7.50±1.118
	p=.241	p=.201	p=.054	p=.896
<b>Education</b>				
Illiterate	7.13±1.251	7.13±.934	7.05±1.022	7.41±1.380
Literate	6.86±1.589	6.65±1.195	6.33±1.507	7.53±1.505
	p=.278	<b>p=.012*</b>	<b>p=.001*</b>	p=.736
<b>Occupation</b>				
Housewife	7.12±1.259	7.10±.950	7.02±1.074	7.45±1.353
Day labor	6.87±1.927	6.83±1.337	6.45±1.167	6.14±2.093
	p=.543	p=.375	p=.104	<b>p=.022*</b>
<b>Type of family</b>				
Nuclear	7.23±1.230	7.17±.985	7.06±1.064	7.39±1.358
Joint	7.04±1.304	7.05±.946	6.97±1.088	7.44±1.416
	p=.163	p=.256	p=.401	p=.786
<b>Number of children</b>				
0-3	7.08±1.313	7.10±1.035	7.00±1.059	7.31±1.372
4 and more	7.14±1.247	7.09±.887	7.00±1.099	7.53±1.399
	p=.677	p=.954	p=.993	p=.230
<b>Number of Family Member</b>				
1-6	7.10±1.295	7.11±.956	6.99±1.095	7.43±1.358
7-12	7.17±1.212	7.03±.987	7.04±1.015	7.40±1.511
	p=.669	p=.509	p=.746	p=.907
<b>Education (Husband)</b>				
Illiterate	7.11±1.274	7.12±.959	7.05±1.032	7.41±
Literate	7.11±1.315	6.92±.967	6.70±1.309	7.49±
	p=.987	p=.170	<b>p=.028*</b>	p=.761
<b>Occupation (Husband)</b>				
Farmer	7.14±1.259	7.07±.986	7.01±1.080	7.45±1.374
Day labor	7.12±1.313	7.25±.855	7.17±.947	7.07±1.709
Businessman	7.09±1.172	6.90±.828	6.67±1.139	7.86±.587
Retired	6.90±1.490	7.07±1.181	6.89±1.275	7.34±1.403
	p=.819	p=.287	p=.116	p=.136
<b>Type of house</b>				
Klay house	7.04±1.327	7.12±.956	7.06±.987	7.52±1.247

Semi Concrete house	7.23±1.185	7.09±1.009	7.07±1.078	7.35±1.429
Concrete house	7.21±1.222	6.97±.892	6.60±1.386	7.11±1.835
	p=.362	p=.590	<b>p=.017*</b>	p=.321
<b>Monthly expenditure</b>				
<500	7.16±1.267	7.01±1.006	6.98±1.035	7.37±1.357
500-1000	6.99±1.379	7.07±.972	6.95±1.216	7.54±1.266
>1000	7.19±1.122	7.30±.809	7.14±.919	7.30±1.693
	p=.429	p=.076	p=.419	p=.584
<b>Monthly income (Family)</b>				
<5000	7.21±1.261	7.07±1.087	7.04±1.042	7.39±1.396
5000-10000	6.95±1.363	7.08±.958	6.95±1.120	7.45±1.292
>10000	7.19±1.191	7.12±.865	7.02±1.068	7.41±1.478
	p=.176	p=.885	p=.778	p=.957
<b>Monthly expenditure(family)</b>				
<5000	7.13±1.389	7.08±1.065	7.01±1.069	7.37±1.387
5000-10000	7.01±1.260	7.06±.956	6.96±1.115	7.40±1.402
>10000	7.20±1.196	7.14±.874	7.04±1.051	7.48±1.385
	p=.464	p=.823	p=.848	p=.888
<b>Age of menopause</b>				
40-45	7.12±1.299	7.06±1.024	7.01±1.076	7.42±1.517
46-50	7.06±1.264	7.07±.906	7.05±.975	7.36±1.278
51-55	7.17±1.276	7.20±.938	6.92±1.240	7.52±1.323
	p=.818	p=.484	p=.676	p=.830

## Discussion

Menopause is a natural transition that all women experience, but individual responses to menopause and the associated drop in estrogen can vary widely due to a range of genetic, cultural, lifestyle, socioeconomic, educational, and dietary influences. In recent years, menopause has gained attention as a significant factor in women's health. In this study, we assessed the quality of life (QOL) in women experiencing menopausal symptoms using the MENQOL (Menopause-Specific Quality of Life) scale. Numerous studies have shown that quality of life (QOL) is often reduced in menopausal women, as this phase is associated with various physical and psychological changes that can affect overall health outcomes [14]. Therefore, QOL of post-menopausal women is needed to be assessed. Thus, in this study, we have tried to evaluate the QOL of post-menopausal women both as a specific and cumulative effect of the four major domains related to her health and well-being, namely vasomotor, physical, sexual and psychosocial.

In the present study, the mean age of menopause was found to be 47.13 ± 4.30 years, which closely aligns with findings from several previous studies conducted in different regions. These include research by Karma et al. in Punjab [14], Kamal and Seedhom in Egypt [23], Waheed et al. in Pakistan [9], Sagdeo and Arora in Nagpur [24], Poomala and Arounassalame in Puducherry [25], Sarkar et al. in Jamnagar [26], and Bansal et al. in Punjab [27]. However, this figure

is lower compared to the study by Nisar and Sohoo in Sindh, Pakistan [18], where the average age of menopause was reported as 52.17 ± 6.019 years.

In the current study shows the most common symptoms reported were decreases in physical strength (100%), feeling lack of energy (100%), changes in appearance, texture or tone of skin (100%), accomplishing less than I used to (99.7%), dry skin (99.7%), decreased stamina (99.7%), sweating (98.2%) and poor memory (98.2%). Whereas Kamal and Seedhom [23] showed that the most frequently reported menopausal symptoms were joint and muscular discomfort (82.1%) followed by physical and mental exhaustion (69.6%) and hot flushes (53.6%).

In the current study, the most severe symptoms identified in the vasomotor, psychosocial, physical, and sexual domains were hot flushes (85.7%), poor memory (92.4%), decrease physical strength (86.2%), and changes in sexual desire (52.6%) respectively. The most severe symptoms of the vasomotor, psychosocial and sexual domain were similar to the study done by Karma et al. in Punjab [14] and Mohamed et al. in Egypt [7]. However, the most severe physical symptoms differed from the present study. In their research, feeling tired or worn out (88%) and low back pain (41.9%) were reported as the most severe physical symptoms.

In this study, the highest mean score in the MENQOL domains was observed in the sexual domain ( $7.42 \pm 1.387$ ), followed by the vasomotor domain ( $7.11 \pm 1.278$ ), then psychosocial domain ( $7.09 \pm 0.961$ ), and lastly, the physical domain ( $7.00 \pm 1.079$ ). These findings align with the study conducted by Mohamed et al. in Egypt [7]. However, the results contradict those of Shobeiri et al. in Iran [4] where the highest mean score of the MENQOL domain was the physical domain ( $39.12 \pm 1.95$ ), followed by the psychosocial domain ( $19.36 \pm 1.20$ ), the vasomotor domain ( $11.65 \pm 5.93$ ), and the sexual domain ( $11.02 \pm 5.66$ ).

In this study, the highest mean scores of symptoms in vasomotor, psychosocial, physical, and sexual domains were hot flushes ( $7.49 \pm 1.101$ ), experiencing poor memory ( $7.66 \pm 0.982$ ), aching muscles and joints ( $7.57 \pm 1.148$ ), and vaginal dryness during intercourse ( $7.57 \pm 1.056$ ), respectively. In contrast, Shobeiri et al. in Iran [4] reported different findings, with the highest mean scores being night sweats ( $4.17 \pm 2.08$ ) for the vasomotor domain, anxiety or nervousness ( $3.34 \pm 2.14$ ) for the psychosocial domain, muscle and joint aches ( $3.41 \pm 2.04$ ) for the physical domain, and changes in sexual desire ( $3.77 \pm 2.11$ ) for the sexual domain.

Previous studies [28] examining the link between menopausal symptoms and various socio-demographic and lifestyle factors found that quality of life (QOL) was associated with lower socio-economic

status, educational level, duration of menopause, physical activity, employment status, and age. In our study, we identified statistically significant associations between several of these factors and specific domains of QOL. Age was linked to the psychosocial domain; educational qualification correlated with both the psychosocial and physical domains; occupation was associated with the sexual domain; the husband's educational qualification was related to the physical domain; and the type of housing was also connected to the physical domain, all with  $p$ -values  $\leq 0.05$ . However, a study by Karma et al. in Punjab [14] reported no statistically significant associations between age, education, occupation, number of children, and various menopausal symptoms.

## Conclusion

The mean scores across each domain indicate that menopausal symptoms were linked to a decline in quality of life among the study participants. Key factors influencing postmenopausal quality of life included age, educational level, occupation, the husband's educational qualifications, and the type of housing. Women require increased care and support during the postmenopausal phase. Hence, developing effective intervention programs is essential to enhance the quality of life for postmenopausal women.

## References

- Moridi G, Shahoei R, Khaldi S, Sayedolshohadaei F (2013) Quality of life among Iranian postmenopausal women participating in a health educational program. *Chronic Dis J*. 1(2): 63-66.
- Inceboz U, Demirci H, Özbasabaran F, Coban A, Nehir S (2010) Factors affecting the quality of life in climacteric women in Manisa Region. *Trakya Univ Tip Fak Derg*. 27(2): 111-116.
- Ray S, Dasgupta A (2012) An assessment of QOL and its determining factors of postmenopausal women in a rural area of West Bengal, India: A multivariate analysis. *Int J Med Public Health*. 2(4): 12-19.
- Shobeiri F, Jenabi E, Hazavehei SM, Roshanaei G (2016) Quality of life in postmenopausal women in Iran: a population-based study. *J Menopausal Med*. 22(1): 31-8.
- Orley J, Kuyken W, editors. *Quality of life assessment: international perspectives*. Berlin, Germany: Springer Berlin Heidelberg.
- Mousavi SA, Masoumi SZ, Keramat A, Pooralajal J, Shobeiri F (2013) Assessment of questionnaires measuring quality of life in infertile couples: a systematic review. *J Reprod Infertil*. 14(3): 110-9.
- Mohamed HA, Lamadah SM, Zamil LGA (2017) Quality of life among menopausal women. *Int J Reprod Contracept Obstet Gynecol*. 3(3): 552-561.
- Greenblum CA, Rowe MA, Neff DF, Greenblum JS (2013) Midlife women: symptoms associated with menopausal transition and early postmenopausal and quality of life. *Menopause*. 20(1): 22-7.
- Waheed K, Khanum A, Butt A, Ejaz S, Randhawa F (2016) Quality of life after menopause in Pakistani women. *Gynecol Obstet (Sunnyvale)*. 6(367): 2161-0932.
- Winblad B, Gauthier S, Astrom D, Stender K (2010) Memantine benefits functional abilities in moderate to severe Alzheimer's disease. *J Nutr Health Aging*. 14(9): 770-4.
- Bashar M, Ahmed K, Uddin MS, Ahmed F, Emran AA, et al. (2017) Depression and quality of life among postmenopausal women in Bangladesh: a cross-sectional study. *J Menopausal Med*. 23(3): 172-181.
- Cheng MH, Lee SJ, Wang SJ, Wang PH, Fuh JL (2007) Does menopausal transition affect the quality of life? A longitudinal study of middle-aged women in Kinmen. *Menopause*. 14(5): 885-90.
- Fuh JL, Wang SJ, Lee SJ, Lu SR, Juang KD (2003) Quality of life and menopause transition for middle-aged women on Kinmen Island. *Qual Life Res*. 12(1): 53-61.
- Karma D, Saini S, Singh S, Sehgal DK (2018) Quality of life among post-menopausal women in rural area of Punjab. *Journal of Medical Science and Clinical Research*. 6(1): 32471-7.



15. Padubidri VG, Daftary SN (2006) Menopause. 13th ed. New Delhi: Reed Elsevier;. Shaw's Textbook of Gynaecology. p. 56-67.
16. Howkins J, Bourne G. Perimenopause, menopause, premature menopause and postmenopausal bleeding. In: Padubidri VG, Daftary SN, editors. Shaw's Textbook of Gynaecology. 14th ed. India: Elsevier; 2008. p. 37.
17. World Health Organization (1996) Research on the menopause in the 1990s. Report of a WHO scientific group. World Health Organization Technical Report Series. 866: 1-107.
18. Nisar N, Sohoo NA (2009) Frequency of menopausal symptoms and their impact on the quality of life of women: A hospital-based survey. *J Pak Med Assoc.* 59(11): 752-6.
19. Sulak PJ (1996) The perimenopause: a critical time in a woman's life. *Int J Fertil Menopausal Stud.* 41(2): 85-9.
20. Williams R, Levine B, Kalilani L, Lewis J, Clark R (2009) Menopause-specific questionnaire assessment in US population-based study shows negative impact on health-related quality of life. *Maturitas.* 62(2): 153-9.
21. Whelan TJ, Goss PE, Ingle JN, Pater JL, Tu D, et al. (2005) Assessment of quality of life in MA.17: a randomized placebo-controlled trial of letrozole after 5 years of tamoxifen in postmenopausal women. *J Clin Oncol.* 23(28): 6931-40.
22. Lewis V (2009) Undertreatment of menopausal symptoms and novel options for comprehensive management. *Curr Med Res Opin.* 25(11): 2689-98.
23. Kamal N, Seedhom A (2017) Quality of life among postmenopausal women in rural Minia, Egypt. *East Mediterr Health J.* 23(8): 527-33.
24. Sagdeo MM, Arora D (2011) Menopausal symptoms: a comparative study in rural and urban women. *JK Sci J Med Educ Res.* 13(1): 23-6.
25. Poomala GK, Arounassalame B (2013) The quality of life during and after menopause among rural women. *J Clin Diagn Res.* 7(1): 135-9.
26. Sarkar A, Pithadia P, Goswami K, Bhavsar S, Makwana NR, et al. (2014) A study on health profile of post-menopausal women in Jamnagar district, Gujarat. *J Res Med Dent Sci.* 2: 25-9.
27. Bansal P, Chaudhary A, Soni RK, Kaushal P (2013) Menopausal problems among rural middle-aged women of Punjab. *Int J Res Health Sci.* 1: 103-9.
28. Abedzadeh Kalarhoudi M, Taebi M, Sadat Z, Saberi F (2011) Assessment of quality of life in menopausal periods: a population study in Kashan, Iran. *Iran Red Crescent Med J.* 13(11): 811-7.